

**CONTROL ID:** 2257030

**TITLE:** The formation of aromatic molecules in heavily obscured post-AGB stars with O-rich masers

**ABSTRACT BODY:**

**Abstract Body:** We report Spitzer/IRS spectra of a small sample of heavily obscured and high mass loss post-AGB stars with O-rich maser emission. IRAM 30 m observations for two post-AGBs are also reported. The Spitzer/IRS spectra show the C-rich character of their circumstellar envelopes (CSE), as revealed by the presence of small hydrocarbon molecules such as  $C_2H_2$ ,  $C_4H_2$ ,  $C_6H_2$ ,  $C_6H_6$ , and HCN. Benzene ( $C_6H_6$ ) is detected toward two sources in our sample, bringing up to three the total number of Galactic post-AGB objects where this molecule has been detected. The available IRAM 30 m data confirm that the central stars are C-rich, despite the presence of O-rich masers, and the presence of high velocity molecular outflows together with extreme mass loss rates ( $10^{-4} M_{\odot} \text{ yr}^{-1}$ ). Our Spitzer and IRAM 30 m observations confirm the model of Cernicharo (2004) that predicts a rich photochemistry in the neutral regions of these objects on timescales shorter than the dynamical evolution of the central HII region, leading to the formation of large, complex carbon-rich molecules. This model also explain the presence of O-rich masers in C-rich post-AGB stars as a consequence of the extreme mass loss rate, being not necessary to invoke a chemistry change in their circumstellar envelopes.

**CONTACT (NAME ONLY):** Pedro Garcia-Lario

**CONTACT (E-MAIL ONLY):** Pedro.Garcia.Lario@esa.int

**AUTHORS/INSTITUTIONS:** P. Garcia-Lario, J. Perea-Calderon, European Space Astronomy Centre, European Space Agency, Madrid, SPAIN|D.A. Garcia-Hernandez, Instituto de Astrofisica de Canarias, La Laguna, Tenerife, SPAIN|J. Cernicharo, ICMM, Madrid, SPAIN|D. Engels, Hamburger Sternwarte, Hamburg, GERMANY|

**PRESENTATION TYPE:** Oral